

Rapid Flex[®] LQ

Complete Liquid Joint Supplement For Horses



~Fast Acting, has been proven in clinical trials to show significant results, fast

~Liquid form = faster absorption

~Clinically proven formula, trials show that Rapid Flex improves range of motion and ability to put weight on affected limbs

~Patented Formula, the only product with this unique joint protection system

~No Devils Claw or Cayenne; ideal for show enthusiasts or actively reproductive broodmares

~Measuring cup included for no-mess use!

Extremely Palatable- Horses love our tasty banana flavor!

Product Information

PI Code (cs):	05-9326-5335
UPC Code (ea):	0-95668-93261-9
Size:	6/32 oz
Package Dimension:	9.6 x 3.25 x 3.25
Species:	Equine
Feed Form:	Liquid

Rapid Flex[®] LQ

Active Ingredients Per Scoop

Glucosamine HCl (shellfish)	5,000 mg
Chondroitin Sulfate (porcine)	850 mg
Hyaluronic Acid	100 mg
Yucca Root	500 mg
Alfalfa	500 mg
Nettle Leaf	100mg
Turmeric Root	50 mg
Celery Seed	25 mg
Black Cohosh Root	25 mg
Ginger Root	25 mg

Inactive Ingredients

Artificial flavor, citric acid, distilled water, methylparaben, potassium sorbate, propylparaben, sucrose, xanthan gum.

Directions For Use

Shake well before each use, Use the measuring cup provided:

Initial Administration (3 weeks)

Average Horses 2 fluid ounces per day
Horses over 1200 lb. 3 fluid ounces per day

Maintenance Administration

Average Horses 1 fluid ounce per day
Horses over 1200 lb. 1 ½ fluid ounces per day

Reminder: Feed is perishable. Store this product in a cool, dry area away from insects and rodents. Do not store below 32° F

Warning: For animal use only. Keep out of reach of children and other animals. In case of accidental overdose, contact a health professional immediately.

Warning: This product should not be given to animals intended for human consumption.

Cautions: Safe use in pregnant animals or animals intended for breeding has not been proven. Consult with your veterinarian before using this product in animals: with clotting disorders; being treated with anticoagulant medications; diabetes, or any metabolic disorder causing hyperglycemia; history of urinary tract stones; known allergies to shellfish. If lameness worsens, discontinue use and contact your veterinarian. Administer during or after the animal has eaten to reduce incidence of gastrointestinal upset.