



Compendium of Veterinary Products - US edition

DURVET, INC.

100 S.E. MAGELLAN DRIVE, BLUE SPRINGS, MO, 64014

Telephone: 816-229-9101

Toll-Free: 800-821-5570

Fax: 816-224-3080

Website: www.durvet.com

Email: info@durvet.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

CANINE SPECTRA[®] 9

Durvet

Canine Distemper-Adenovirus Type 2-Parainfluenza-Parvovirus Vaccine

Modified Live Virus

Leptospira Canicola-Grippotyphosa-Icterohaemorrhagiae-Pomona Bacterial Extract

FOR DOGS

KEEP PRODUCT REFRIGERATED

Description: Spectra[®] 9 vaccine is a combination of immunogenic, attenuated strains of Canine Distemper, Canine Adenovirus Type 2 (CAV-2), Canine Parainfluenza, and Canine Parvovirus Type 2b, propagated in cell line tissue cultures. The diluent contains killed antigens for *Leptospira Canicola-Grippotyphosa-Icterohaemorrhagiae-Pomona* bacterial extract. The CAV-2 fraction cross-protects against respiratory infection caused by infectious canine hepatitis (CAV-1).

Indications: For subcutaneous vaccination of healthy dogs 6 weeks of age or older as an aid in the prevention of disease caused by canine distemper, infectious canine hepatitis, canine adenovirus type 2, canine parainfluenza, canine parvovirus (CPV), *Leptospira canicola*, *Leptospira grippotyphosa*, *Leptospira icterohaemorrhagiae*, and *Leptospira pomona*. This product contains a CPV 2b strain.

Directions and Dosage: The dosage is 1 mL injected subcutaneously. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe. Thoroughly prepare injection site with antiseptic. Lift the loose skin behind the neck or behind front leg and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel.

IMPORTANT NOTE: *Before injecting vaccine, pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site.* A recommended vaccination schedule should start at or about 6 weeks of age. The presence of maternal antibody is known to interfere with the development of active immunity. Puppies should be revaccinated every 2 to 3 weeks until at least 16 weeks of age. All dogs over 12 weeks of age should initially receive one dose of Spectra[®] 9 and a second dose 2 to 3 weeks later. Annual revaccination with a single dose is recommended.



Precautions: Only vaccinate healthy animals. Animals incubating any disease or stressed due to shipping, malnutrition, or parasitism may not achieve or maintain an adequate immune response. Burn containers and unused contents. Although rare, severe allergic reactions (anaphylactoid) may occur that require immediate veterinary care.

Antidote: Epinephrine.

Storage: Store out of direct sunlight at 35°-45°F (2°-7°C). Do not freeze.

Preservatives: Gentamicin and thimerosal.

Produced by: Boehringer Ingelheim Vetmedica, Inc., St. Joseph, MO 64506 U.S.A.

U.S. Vet. License No. 124

Distributed by: Peak Marketing, P.O. Box 279, Blue Springs, MO 64013 USA

Spectra[®] 9 is a trademark of Peak Marketing LLC

MADE IN THE USA

One Dose with Syringe	Package contains a one dose vial of vaccine, a one mL vial of vaccine diluent, & one disposable syringe	151211-00	ISS13XB09
25 Single Doses with Syringe			ISS13XB09

CPN: 1084355.0

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated March 23, 2018

Copyright 2018 - Animalytix LLC